



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

$$\begin{aligned}
\left(1 - \frac{4x^2}{\pi^2}\right)\left(1 - \frac{4x^2}{3^2\pi^2}\right) \cdot \dots = 1 - \frac{4x^2}{\pi^2} \left[\frac{1}{1^2} + \frac{1}{3^2} + \frac{1}{5^2} + \dots \right] \\
+ \frac{4^2x^4}{\pi^4} \left[\frac{1}{3^2} + \frac{1}{3^2 \cdot 5^2} + \dots \right] - \frac{4^3x^6}{\pi^6} \left[\frac{1}{3^2 \cdot 5^2} + \dots \right] + \dots \\
= 1 - c_1x^2 + c_2x^4 - c_3x^6 + \dots \quad (3).
\end{aligned}$$

Comparing (1), (2), and (3),

$$\begin{aligned}
1 - c_1x^2 + c_2x^4 - c_3x^6 \cdot \dots = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \frac{x^6}{6!} + \dots \\
\text{or } \left(1 - \frac{4x^2}{\pi^2}\right)\left(1 - \frac{4x^2}{3^2\pi^2}\right) \cdot \dots = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} \cdot \dots \quad (4).
\end{aligned}$$

$$\text{Let } y^2 = -x^2. \quad \left(1 + \frac{4y^2}{\pi^2}\right)\left(1 + \frac{4y^2}{3^2\pi^2}\right) \cdot \dots = 1 + \frac{y^2}{2!} + \frac{y^4}{4!} + \frac{y^6}{6!} + \dots \quad (5).$$

Now, adding this exponential series, e^y and e^{-y} ,

$$\frac{e^y + e^{-y}}{2} = 1 + \frac{y^2}{2!} + \frac{y^4}{4!} + \dots = \text{from (5)} \quad \left(1 + \frac{4y^2}{\pi^2}\right)\left(1 + \frac{4y^2}{3^2\pi^2}\right) \cdot \dots \quad (6).$$

Substituting π for y in (6), we have,

$$e^\pi + e^{-\pi} = 2[1 + 2^2][1 + (\frac{2}{3})^2][1 + (\frac{2}{5})^2] \cdot \dots$$

PROBLEMS FOR SOLUTION.

ARITHMETIC.

132. Proposed by WILLIAM SYMMONDS, A.M., Professor of Mathematics, Santa Rosa College, Sebastopol, Cal.

A road 60 feet wide crosses a square acre of land. The west line of the road passes through the southwest corner of the land, while the east line of the former passes through the northeast corner of the latter. What fraction of the land is included in the road?

133. Proposed by COOPER D. SCHMITT, A. M., Professor of Mathematics, University of Tennessee. Knoxville, Tenn.

In Wentworth's Arithmetic he gives a formula $\frac{\pi}{4} \frac{1}{d} (d^2 - 2d)$ for calculating the number of board feet in a log 10 feet long, when d is the diameter in inches. How is this rule derived?

*** Solutions of these problems should be sent to B. F. Finkel not later than Nov. 10.